

ON THE ISOMETRIC COMPOSITION OPERATORS ON THE BLOCH SPACE IN $\mathbb{C}^n$

ROBERT F. ALLEN AND FLAVIA COLONNA

ABSTRACT. Let φ be a holomorphic self-map of a bounded homogeneous domain D in $\mathbb{C}^n$. In this work, we show that the composition operator $C_\varphi : f \mapsto f \circ \varphi$ is bounded on the Bloch space $\mathcal{B}$ of the domain and provide estimates on its operator norm. We also give a sufficient condition for φ to induce an isometry on $\mathcal{B}$. This condition allows us to construct non-trivial examples of isometric composition operators in the case when D has the unit disk as a factor. We then obtain some necessary conditions for C_φ to be an isometry on $\mathcal{B}$ when D is a Cartan classical domain. Finally, we give the complete description of the spectrum of the isometric composition operators in the case of the unit disk and for a wide class of symbols on the polydisk.

1. INTRODUCTION

An analytic function f on $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ is said to be **Bloch** if

$$(1) \quad \beta_f = \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| < \infty.$$

The set $\mathcal{B}$ of Bloch functions on $\mathbb{D}$ is a Banach space with semi-norm $f \mapsto \beta_f$ and norm $\|f\|_{\mathcal{B}} = |f(0)| + \beta_f$.

By the Schwarz-Pick lemma, if f is any Bloch function on $\mathbb{D}$ and φ is an analytic function mapping $\mathbb{D}$ into itself, then $f \circ \varphi \in \mathcal{B}$ and $\beta_{f \circ \varphi} \leq \beta_f$, with equality holding if $\varphi \in \text{Aut}(\mathbb{D})$, where $\text{Aut}(D)$ denotes the set of biholomorphic transformations of a domain D (which we call **automorphisms** of D).

In (30), Xiong proved that the composition operator $C_\varphi(f) = f \circ \varphi$ is bounded on $\mathcal{B}$ and gave estimates for its operator norm. Furthermore, he obtained several necessary conditions for C_φ to be an isometry (that is, to preserve the Bloch norm). The function φ is called the **symbol** of the operator C_φ .

In (7), the second author completely characterized the symbols of the isometric composition operators. In Theorem 1.1, we list several equivalent conditions that can be used to describe the class of such symbols.

Theorem 1.1. *Let φ be an analytic function mapping $\mathbb{D}$ into itself. Then C_φ is an isometry on $\mathcal{B}$ if and only if $\varphi(0) = 0$ and any of the following equivalent conditions holds:*

- (a) $\beta_\varphi = 1$.
- (b) $B_\varphi := \sup_{z \in \mathbb{D}} \frac{(1 - |z|^2) |f'(z)|}{1 - |f(z)|^2} = 1$.

2000 *Mathematics Subject Classification.* primary: 30D45, secondary: 47B38, 47A30.

Key words and phrases. Composition operators, Bloch space, Homogeneous domains, Isometry.

(c) Either $\varphi \in \text{Aut}(\mathbb{D})$ or for every $a \in \mathbb{D}$ there exists a sequence $\{z_k\}$ in $\mathbb{D}$ such that $|z_k| \rightarrow 1$, $\varphi(z_k) = a$, and

$$\lim_{k \rightarrow \infty} \frac{(1 - |z_k|^2)|\varphi'(z_k)|}{1 - |\varphi(z_k)|^2} = 1.$$

(d) Either $\varphi \in \text{Aut}(\mathbb{D})$ or for every $a \in \mathbb{D}$ there exists a sequence $\{z_k\}$ in $\mathbb{D}$ such that $|z_k| \rightarrow 1$, $\varphi(z_k) \rightarrow a$, and

$$\lim_{k \rightarrow \infty} \frac{(1 - |z_k|^2)|\varphi'(z_k)|}{1 - |\varphi(z_k)|^2} = 1.$$

(e) Either $\varphi \in \text{Aut}(\mathbb{D})$ or the zeros of φ form an infinite sequence $\{z_k\}$ in $\mathbb{D}$ such that

$$\limsup_{k \rightarrow \infty} (1 - |z_k|^2)|\varphi'(z_k)| = 1.$$

(f) Either $\varphi \in \text{Aut}(\mathbb{D})$ or $\varphi = gB$, where g is a non-vanishing analytic function mapping $\mathbb{D}$ into itself or a constant of modulus 1, and B is an infinite Blaschke product whose zero set Z contains a sequence $\{z_k\}$ such that $|g(z_k)| \rightarrow 1$ and

$$\lim_{k \rightarrow \infty} \prod_{\zeta \in Z, \zeta \neq z_k} \left| \frac{z_k - \zeta}{1 - \overline{z_k}\zeta} \right| = 1.$$

(g) Either $\varphi \in \text{Aut}(\mathbb{D})$ or there exists $\{S_k\}_{k \in \mathbb{N}}$ in $\text{Aut}(\mathbb{D})$ such that $|S_k(0)| \rightarrow 1$ and $\{\varphi \circ S_k\}$ approaches the identity locally uniformly in $\mathbb{D}$.

Condition (d) was noted in (20). For the equivalence of the other conditions see (7), (5) and (6).

Condition (f) yields a recipe for constructing examples of isometric composition operators induced by symbols that are not rotations. Easily constructible examples are Blaschke products whose zero sets $\{z_k\}$ are **thin**, that is, satisfy the condition

$$\lim_{k \rightarrow \infty} \prod_{j \neq k} \left| \frac{z_k - z_j}{1 - \overline{z_k}z_j} \right| = 1.$$

In this article, we study the symbols of the isometric composition operators on the Bloch space of a bounded homogeneous domain in $\mathbb{C}^n$ ($n \in \mathbb{N}$, $n \geq 2$) and, when needed, restrict our attention to bounded symmetric domains. Our goal is obtain in higher dimensions conditions analogous to (a)-(g) in Theorem 1.1 to whatever extent is possible.

1.1. Organization of the paper. In Section 2, we review the background on Bloch functions on a bounded homogeneous domain in $\mathbb{C}^n$. We also recall the Carthan classification of bounded symmetric domains and the corresponding invariant metrics, known as **Bergman metrics**.

In Section 3, we give a characterization of the Bloch semi-norm of a Bloch function on a bounded homogeneous domain as the maximum dilation of f with respect to the Bergman distance, which allows us to show that the composition operator C_φ is bounded on the Bloch space $\mathcal{B}$ of the domain. We also provide estimates for its operator norm. The boundedness of C_φ was observed in (25) and in (29), but the authors only showed boundedness with respect to the Bloch semi-norm (which yields the boundedness with respect to the Bloch norm under the assumption that φ fixes the base point of the domain used to define the Bloch norm). We then show that the correspondence $f \mapsto \beta_f$ is lower-semi-continuous on the Bloch space.

In Section 4, we recall the classification of the Bloch constant of a bounded symmetric domain D (namely, the supremum of the Bloch semi-norms of the holomorphic functions mapping D into $\mathbb{D}$) in terms of the Cartan class and the dimension of D . We then show that for most domains, the canonical projections map D into $\mathbb{D}$ and have maximal Bloch semi-norm, a property that will be used in Section 6 to derive a necessary condition for a holomorphic self-map of a bounded symmetric domain of this type to induce an isometry on the Bloch space.

In Section 5, we give some simple properties of the maximum Bergman dilation of a holomorphic self-map and use the lower-semi-continuity of the Bloch semi-norm to obtain a sufficient condition for a holomorphic self-map on a bounded homogeneous domain D to induce an isometry on $\mathcal{B}$ under composition. We use this result to generate non-trivial examples in the special case when D has $\mathbb{D}$ as a factor.

In Section 6, we give some necessary conditions for a holomorphic self-map of D to be the symbol of an isometric composition operator on $\mathcal{B}$ when D is a product of Cartan classical domains. We conclude the section with a conjecture.

In Section 7 we analyze the spectrum of C_φ when this is an isometry on $\mathcal{B}$ in the case when D is the unit polydisk. We deduce the full description of the spectrum in the one-dimensional case.

We conclude the paper (Section 8) with a list of open questions.

2. BLOCH FUNCTIONS IN HIGHER DIMENSIONS

A **homogeneous domain** in $\mathbb{C}^n$ is a domain D such that $\text{Aut}(D)$ acts transitively on D , i.e. given any two points $z, w \in D$ there exists $T \in \text{Aut}(D)$ such that $T(z) = w$.

In (14), Hahn introduced the notion of Bloch function on a bounded homogeneous domain in $\mathbb{C}^n$.

Let f be a complex-valued holomorphic function on a bounded homogeneous

domain D in $\mathbb{C}^n$. For $u, v \in \mathbb{C}^n$, let $\langle u, v \rangle = \sum_{k=1}^n u_k \overline{v_k}$, and for $z \in D$, let

$(\nabla f)(z)u = \langle (\nabla f)(z), \overline{u} \rangle$, where $(\nabla f)(z)$ is the gradient of f at z . Denote by $H_z(u, \overline{u})$ the Bergman metric on D , that is, the positive definite Hermitian form which is invariant under automorphisms of D . Using a different approach from Hahn's, in (23) Timoney defined a Bloch function on D as a holomorphic function f on D such that $\beta_f = \sup_{z \in D} Q_f(z)$ is finite, where

$$Q_f(z) = \sup_{u \in \mathbb{C}^n \setminus \{0\}} \frac{|(\nabla f)(z)u|}{H_z(u, \overline{u})^{1/2}}.$$

Denote by $\mathcal{B}$ the space of Bloch functions on D . The map $f \mapsto \beta_f$ is a semi-norm on $\mathcal{B}$, and fixing any point $z_0 \in D$, the set $\mathcal{B}$ is a Banach space, called the **Bloch space**, under the norm $\|f\|_{\mathcal{B}} = |f(z_0)| + \beta_f$. For convenience, throughout this paper we shall assume that $0 \in D$ and $z_0 = 0$. The Bloch space contains the space of bounded holomorphic functions (23). When $D = \mathbb{D}$, the above definition of Bloch function reduces to condition (1).

Excellent references on Bloch functions include (1) for the one-variable case, (23) and (24) for the several-variable case. See also (3) for an overview of Bloch functions on the disk and its connection to other function spaces. Bloch functions have

been defined on more general classes of bounded domains, such as strongly pseudoconvex domains (19). These domains, however, have sparse, and often trivial, automorphism groups, and operator theory problems are much more difficult to treat.

A domain D in $\mathbb{C}^n$ is said to be **symmetric** if for each $z_0 \in D$, there exists an automorphism S of D which is an involution (i.e. $S \circ S$ is the identity) and has z_0 as an isolated fixed point. Symmetric domains are homogeneous (see (14), pp. 170, 301). Examples of symmetric domains are the unit ball

$$\mathbb{B}_n = \{z = (z_1, \dots, z_n) \in \mathbb{C}^n : \|z\| < 1\},$$

where $\|z\| = (\sum_{k=1}^n |z_k|^2)^{1/2}$, and the unit polydisk

$$\mathbb{D}^n = \{z = (z_1, \dots, z_n) \in \mathbb{C}^n : |z_k| < 1 \text{ for all } k = 1, \dots, n\}.$$

We now describe their groups of automorphisms, and their Bergman metrics.

The automorphism group of the unit ball can be described as

$$\text{Aut}(\mathbb{B}_n) = \{U \circ \varphi_a : U \text{ unitary}, a \in \mathbb{B}_n\},$$

where for $z \in \mathbb{B}_n$, $\varphi_0(z) = z$, while for $a \in \mathbb{B}_n \setminus \{0\}$,

$$(2) \quad \varphi_a(z) = \frac{a - P_a(z) - (1 - \|a\|^2)^{1/2} Q_a(z)}{1 - \langle z, a \rangle},$$

with $P_a(z) = \frac{\langle z, a \rangle}{\|a\|^2} a$, and $Q_a(z) = z - P_a(z)$ (see (21), Theorem 2.2.5). For $z \in \mathbb{B}_n$, $u, v \in \mathbb{C}^n$, the Bergman metric on $\mathbb{B}_n$ is given by

$$H_z(u, \bar{v}) = \frac{(n+1)[(1 - \|z\|^2)\langle u, v \rangle + \langle u, z \rangle \langle z, v \rangle]}{2(1 - \|z\|^2)^2}.$$

The automorphism group of the polydisk is given by

$$\text{Aut}(\mathbb{D}^n) = \{(T_1(z_{\tau(1)}), \dots, T_n(z_{\tau(n)})) : T_k \in \text{Aut}(\mathbb{D}), \tau \in S_n\},$$

where S_n is the group of permutations on the set $\{1, \dots, n\}$ (see (22), p. 167). For $z \in \mathbb{D}^n$, $u, v \in \mathbb{C}^n$, the Bergman metric on $\mathbb{D}^n$ is given by

$$H_z(u, \bar{v}) = \sum_{k=1}^n \frac{u_k \bar{v}_k}{(1 - |z_k|^2)^2}.$$

For a function φ mapping a domain $D \subset \mathbb{C}^n$ into $\mathbb{C}^n$ and for $z \in D$, let $J\varphi(z)$ be the Jacobian matrix of φ at z , that is, the $n \times n$ matrix whose (j, k) -entry is $\partial \varphi_j(z) / \partial z_k$. By definition of the Bergman metric, if D is a bounded homogeneous domain and $S \in \text{Aut}(D)$, then for all $z \in D$, and for all $u \in \mathbb{C}^n$

$$(3) \quad H_{S(z)}(JS(z)u, \overline{JS(z)u}) = H_z(u, \bar{u}),$$

where $JS(z)u$ is the usual matrix product where u is viewed as a column vector. As an immediate consequence of (3), we deduce that the Bloch semi-norm is invariant under right composition of automorphisms.

We use the notation $H(D, D')$ for the class of holomorphic functions mapping a domain D in $\mathbb{C}^n$ into a domain D' in $\mathbb{C}^m$ and use the abbreviation $H(D)$ for $H(D, D)$.

In (23) (proof of Theorem 2.12) it was shown that for $n, m \in \mathbb{N}$, if $D \subset \mathbb{C}^n$ and $D' \subset \mathbb{C}^m$ are bounded homogeneous domains, then there is a constant $c > 0$ depending only on D and D' such that for any $\varphi \in H(D, D')$

$$H_{\varphi(z)}^{D'}(J\varphi(z)u, \overline{J\varphi(z)u}) \leq cH_z^D(u, \overline{u}),$$

where H_z^D and $H_{\varphi(z)}^{D'}$ are the Bergman metrics on D and D' at z and $\varphi(z)$, respectively. In particular, if D is a bounded homogeneous domain and φ is a holomorphic map of D into itself, then

$$B_\varphi = \sup_{z \in D} \sup_{u \in \mathbb{C}^n \setminus \{0\}} \frac{H_{\varphi(z)}(J\varphi(z)u, \overline{J\varphi(z)u})^{1/2}}{H_z(u, \overline{u})^{1/2}}$$

is bounded above by a constant independent of φ . Thus, for any Bloch function f on D and any $z \in D$, we have

$$Q_{f \circ \varphi}(z) \leq \sup_{u \in \mathbb{C}^n \setminus \{0\}} \left(\frac{H_{\varphi(z)}(J\varphi(z)u, \overline{J\varphi(z)u})}{H_z(u, \overline{u})} \right)^{1/2} Q_f(\varphi(z)).$$

Thus

$$(4) \quad Q_{f \circ \varphi}(z) \leq B_\varphi Q_f(\varphi(z)).$$

Furthermore, $Q_{f \circ \varphi}(z) = Q_f(\varphi(z))$ if $\varphi \in \text{Aut}(D)$. Consequently,

$$(5) \quad \beta_{f \circ \varphi} \leq B_\varphi \beta_f,$$

and so composition operators whose symbol fixes 0 are bounded on $\mathcal{B}$ and the composition operators induced by the automorphisms of D preserve the Bloch semi-norm. Moreover, if C_φ preserves the Bloch semi-norm, then $B_\varphi \geq 1$. In Section 3, we shall prove that C_φ is in fact bounded on the Bloch space without the assumption $\varphi(0) = 0$.

We call B_φ the **Bergman constant** of φ . By (3), the Bergman constant of an automorphism is 1. When $n = 1$, $D = \mathbb{D}$, and φ is any analytic self map of $\mathbb{D}$ we see that $H_z(u, \overline{u}) = \frac{|u|^2}{(1-|z|^2)^2}$. Thus, B_φ is the constant in part (b) of Theorem 1.1.

The one-dimensional case suggests that the value of the Bergman constant may play a role in helping us determine which holomorphic self-maps of a bounded homogeneous domain induce composition operators that are isometries on the Bloch space. The higher-dimensional case is more difficult to treat because, in general, there may exist functions $\varphi \in H(D)$ whose Bergman constant is larger than 1. Indeed, in (17) Korányi proved that the value of the maximal Bergman constant is related to the rank of the domain. We state his result in Theorem 5.2. An example of a function φ whose Bergman constant is larger than 1 is $\varphi(z) = (z_1, \dots, z_1)$ for $z \in \mathbb{D}^n$. However, the corresponding operator C_φ is not an isometry on $\mathcal{B}$. Indeed, as we shall observe in Theorem 6.1, if C_φ is an isometry on $\mathcal{B}$, then the components of φ are linearly independent.

Because of the invariance of the semi-norm under composition by automorphisms, easy examples of symbols of isometric composition operators are the automorphisms that fix 0. In the case of the unit polydisk they are the functions $(z_1, \dots, z_n) \mapsto (\lambda_1 z_{\tau(1)}, \dots, \lambda_n z_{\tau(n)})$ where the λ_j ($j = 1, \dots, n$) are unimodular constants and $\tau \in S_n$. In the case of the unit ball, they are the unitary transformations.

In (6) the problem of characterizing the holomorphic self-maps of the unit poly-disk which are symbols of isometric composition operators was studied. Although a complete characterization was not achieved, several necessary conditions were obtained and a large class of symbols of isometric composition operators was described. Other references on composition operators on the Bloch space in higher dimensions include (25) and (29).

E. Cartan (2) proved that any bounded symmetric domain D is biholomorphically equivalent to a finite product of irreducible bounded symmetric domains, unique up to rearrangement of the factors. He then classified all the irreducible domains into six classes, four of which form infinite families (known as **Cartan classical domains**) and two classes each containing a single domain of dimension 16 and 27, respectively, called **exceptional domains**. All these classes contain the origin. A bounded symmetric domain D is said to be in **standard form** if it has the form $D = D_1 \times \cdots \times D_k$, where each D_j is a Cartan classical domain or an exceptional domain.

The Cartan classical domains $R_I, R_{II}, R_{III}, R_{IV}$ and the corresponding Bergman metrics are described below. The notation we use conforms to Kobayashi's in (16), except we scale the Bergman metrics by dividing them by 4. For a description of their automorphism groups see (16). For a description of the exceptional domains R_V and R_{VI} , see (12).

Let $\mathcal{M}_{m \times n}$ denote the set of $m \times n$ matrices over $\mathbb{C}$ and let $\mathcal{M}_n = \mathcal{M}_{n \times n}$. Let $I_n \in \mathcal{M}_n$ be the identity matrix and let Z^* be the adjoint of the matrix Z . Then

$$\begin{aligned} R_I &= \{Z \in \mathcal{M}_{m \times n} : I_m - ZZ^* > 0\}, \text{ for } m \geq n \geq 1, \\ H_Z(U, \bar{V}) &= \frac{m+n}{2} \text{Trace}[(I_m - ZZ^*)^{-1} U (I_n - Z^* Z)^{-1} V^*], \\ R_{II} &= \{Z \in \mathcal{M}_n : Z = Z^T, I_n - ZZ^* > 0\}, \text{ for } n \geq 1, \\ H_Z(U, \bar{V}) &= \frac{n+1}{2} \text{Trace}[(I_n - ZZ^*)^{-1} U (I_n - Z^* Z)^{-1} V^*], \\ R_{III} &= \{Z \in \mathcal{M}_n : Z = -Z^T, I_n - ZZ^* > 0\}, \text{ for } n \geq 2, \\ H_Z(U, \bar{V}) &= \frac{n-1}{2} \text{Trace}[(I_n - ZZ^*)^{-1} U (I_n - Z^* Z)^{-1} V^*], \\ R_{IV} &= \left\{ z \in \mathbb{C}^n : \left| \sum z_j^2 \right|^2 + 1 - 2\|z\|^2 > 0, \left| \sum z_j^2 \right|^2 < 1 \right\}, 1 \leq n \neq 2, \\ H_z(u, \bar{v}) &= nAu[A(I_n - z^T \bar{z}) + (I_n - z^T \bar{z})Z^* z(I_n - z^T \bar{z})]v^*, \end{aligned}$$

where z^T is the transpose of z and $A = \left| \sum_{j=1}^n z_j^2 \right|^2 + 1 - 2\|z\|^2$.

Since some irreducible domains may belong to different classes, we add the following dimensional restrictions that yield uniqueness: $n \geq 2$ for domains in R_{II} , and $n \geq 5$ for domains in R_{III} and R_{IV} .

3. BLOCH SEMI-NORM AS A LIPSCHITZ NUMBER

In this section, we give an alternate description of the Bloch semi-norm and analyze some useful properties of Bloch functions.

We begin by observing that the Bloch functions on a bounded homogeneous domain D are precisely the Lipschitz maps between D under the distance ρ induced by the Bergman metric and the complex plane under the Euclidean distance. To see this, we recall a result of (13) connecting local derivatives to Lipschitz mappings.

If we consider $f : D \rightarrow \mathbb{C}$ as a map between Riemannian manifolds, then $Q_f(z)$ is exactly the operator norm of $df(z)$, the induced linear transformation on the tangent space at $z \in D$, with respect to the Bergman metric H_z on D and the Euclidean metric on $\mathbb{C}$. Thus, β_f is the supremum of $\|df(z)\|$ over all $z \in D$. The dilation $\text{dil}(f)$ of f is the global Lipschitz number of f :

$$\text{dil}(f) = \sup_{z, w \in D, z \neq w} \frac{|f(z) - f(w)|}{\rho(z, w)}.$$

A **length space** is a Riemannian manifold in which the distance between two points is the infimum of the lengths of geodesics connecting the points. Examples include bounded homogeneous domains under the Bergman metric. An example of a space which is not a length space is the complement of a closed ball in $\mathbb{C}^n$ under the Euclidean metric. In Property 1.8 bis of (13), it is shown that in the special case when M is a length space and $f : M \rightarrow N$ is a smooth map between Riemannian manifolds,

$$\text{dil}(f) = \sup_{x \in M} \|df(x)\|.$$

We deduce

Theorem 3.1. *Let D be a bounded homogeneous domain and let $f \in H(D, \mathbb{C})$. Then f is Bloch if and only if f is a Lipschitz map as a function from D under the Bergman metric and the complex plane under the Euclidean metric. Furthermore*

$$\beta_f = \sup_{z \neq w} \frac{|f(z) - f(w)|}{\rho(z, w)}.$$

A proof that does not use differential geometry can be found in (8) (Theorem 10) and in (27) (Theorem 5.1.6) for the case of the unit disk, and in (28) (Theorem 3.6) for the case of the unit ball in $\mathbb{C}^n$.

The following result is an extension of Theorem 2 of (30) to any bounded homogeneous domain.

Theorem 3.2. *Let D be a bounded homogeneous domain and let $\varphi \in H(D)$. Then C_φ is a bounded operator on the Bloch space of D . Furthermore*

$$1 \leq \|C_\varphi\| \leq \max\{1, \rho(\varphi(0), 0) + B_\varphi\}.$$

In particular, if $\varphi(0) = 0$, then $1 \leq \|C_\varphi\| \leq \max\{1, B_\varphi\}$.

Proof. By Theorem 3.1, for any $f \in \mathcal{B}$, $|f(\varphi(0)) - f(0)| \leq \rho(\varphi(0), 0)\beta_f$, whence

$$|f(\varphi(0))| \leq |f(0)| + |f(\varphi(0)) - f(0)| \leq \|f\|_{\mathcal{B}} + (\rho(\varphi(0), 0) - 1)\beta_f.$$

Thus, using (5) we obtain

$$\|f \circ \varphi\|_{\mathcal{B}} = |f(\varphi(0))| + \beta_{f \circ \varphi} \leq \|f\|_{\mathcal{B}} + (\rho(\varphi(0), 0) - 1 + B_\varphi)\beta_f.$$

If $\rho(\varphi(0), 0) - 1 + B_\varphi \leq 0$, then $\|f \circ \varphi\|_{\mathcal{B}} \leq \|f\|_{\mathcal{B}}$. If $\rho(\varphi(0), 0) - 1 + B_\varphi \geq 0$, then

$$\|f \circ \varphi\|_{\mathcal{B}} \leq (\rho(\varphi(0), 0) + B_\varphi)\|f\|_{\mathcal{B}}.$$

Hence C_φ is bounded on $\mathcal{B}$ and

$$\|C_\varphi\| = \sup_{\|f\|_{\mathcal{B}}=1} \|f \circ \varphi\|_{\mathcal{B}} \leq \max\{1, \rho(\varphi(0), 0) + B_\varphi\}.$$

The lower estimate can be deduced immediately by taking as a test function f the constant function 1. $\square$

The following result is the extension to the unit ball of Corollary 1 and Theorem 2 in (30).

Corollary 3.1. For any $\varphi \in H(\mathbb{B}_n)$,

$$\max \left\{ 1, \frac{1}{2} \log \frac{1 + \|\varphi(0)\|}{1 - \|\varphi(0)\|} \right\} \leq \|C_\varphi\| \leq \max \left\{ 1, \frac{1}{2} \log \frac{1 + \|\varphi(0)\|}{1 - \|\varphi(0)\|} + B_\varphi \right\}.$$

In particular, if $\varphi(0) = 0$, then $\|C_\varphi\| = 1$.

For the proof we need the following result which is part of Theorem 3.1 of (28).

Lemma 3.1. (28) For all $f \in H(\mathbb{B}_n, \mathbb{C})$, $z \in \mathbb{B}_n$,

$$(6) \quad Q_f(z) = [(1 - \|z\|^2)(\|(\nabla f)(z)\|^2 - |Rf(z)|^2)]^{1/2},$$

where $Rf(z) = \sum_{k=1}^n z_k \frac{\partial f}{\partial z_k}(z)$. Therefore

$$\beta_f = \sup_{z \in \mathbb{B}_n} [(1 - \|z\|^2)(\|(\nabla f)(z)\|^2 - |Rf(z)|^2)]^{1/2}.$$

Proof. The upper estimate follows immediately from Theorem 3.2, since by Proposition 1.21 of (28),

$$\rho(\varphi(0), 0) = \frac{1}{2} \log \frac{1 + \|\varphi(0)\|}{1 - \|\varphi(0)\|}.$$

To prove the lower estimate, note that if $\varphi(0) = 0$, then the result follows from Theorem 3.2. So assume $\varphi(0) \neq 0$. For $z \in \mathbb{B}_n$, define

$$f(z) = \frac{1}{2} \log \frac{\|\varphi(0)\| + \langle z, \varphi(0) \rangle}{\|\varphi(0)\| - \langle z, \varphi(0) \rangle}.$$

Then f is holomorphic on $\mathbb{B}$, $f(0) = 0$, and $f(\varphi(0)) = \frac{1}{2} \log \frac{1 + \|\varphi(0)\|}{1 - \|\varphi(0)\|}$. Moreover, using the triangle inequality, we obtain

$$\begin{aligned} \|(\nabla f)(z)\|^2 - |Rf(z)|^2 &= \frac{\|\varphi(0)\|^2(\|\varphi(0)\|^2 - |\langle z, \varphi(0) \rangle|^2)}{|\|\varphi(0)\|^2 - \langle z, \varphi(0) \rangle|^2} \\ &\leq \frac{\|\varphi(0)\|^2}{|\|\varphi(0)\|^2 - \langle z, \varphi(0) \rangle|^2}. \end{aligned}$$

Thus, by (6), we deduce

$$Q_f(z)^2 \leq \frac{(1 - \|z\|^2)\|\varphi(0)\|^2}{|\|\varphi(0)\|^2 - \langle z, \varphi(0) \rangle|^2} = \frac{1 - \|z\|^2}{|1 - \langle z, u_0 \rangle|^2} \leq 1,$$

where $u_0 = \frac{\varphi(0)}{\|\varphi(0)\|}$. On the other hand, it is immediate to check that $Q_f(0) = 1$. Therefore, $f \in \mathcal{B}$ and $\beta_f = 1 = \|f\|_{\mathcal{B}}$. Hence $\|C_\varphi\| \geq \|f \circ \varphi\|_{\mathcal{B}} \geq |f(\varphi(0))|$. Since $\|C_\varphi\| \geq 1$, the lower estimate follows at once. In Section 5 (see paragraph after Theorem 5.2), it will be shown that $B_\varphi \leq 1$. Thus, in the case when $\varphi(0) = 0$, we deduce that $\|C_\varphi\| = 1$. $\square$

The characterization of the Bloch semi-norm in Theorem 3.1 leads to the following convergence theorem.

Theorem 3.3. *Let $\{f_n\}$ be a sequence of Bloch functions on a bounded homogeneous domain D which converges locally uniformly in D to some holomorphic function f . If the sequence $\{\beta_{f_n}\}$ is bounded, then f is Bloch and*

$$\beta_f \leq \liminf_{n \rightarrow \infty} \beta_{f_n}.$$

That is, the function $f \mapsto \beta_f$ is lower semi-continuous on $\mathcal{B}$ under the topology of uniform convergence on compact subsets of D .

Proof. Since $\{\beta_{f_n}\}$ is bounded, then $C = \liminf_{n \rightarrow \infty} \beta_{f_n}$ exists and is non-negative. Also, there exists a subsequence $\{\beta_{f_{n_k}}\}$ which converges to C . If $C = 0$, then f is constant and the result follows at once. So, we assume $C > 0$.

Let $z, w \in D$ and fix $\epsilon > 0$. Choose a positive integer ν so that $|f_{n_k}(z) - f(z)| < \epsilon/2$, $|f_{n_k}(w) - f(w)| < \epsilon/2$, and $\beta_{f_{n_k}} < C + \epsilon$, for all $k \geq \nu$. Then

$$|f(z) - f(w)| < \epsilon + \beta_{f_{n_k}} \rho(z, w) < \epsilon(1 + \rho(z, w)) + C\rho(z, w).$$

Letting $\epsilon \rightarrow 0$, we obtain $|f(z) - f(w)| \leq C\rho(z, w)$. Using Theorem 3.1, we deduce that f is Bloch and $\beta_f \leq C$. $\square$

4. BLOCH CONSTANT OF A BOUNDED SYMMETRIC DOMAIN

For a bounded symmetric domain D define the **Bloch constant** of D as

$$c_D = \sup\{\beta_f : f \in H(D, \mathbb{D})\}$$

and the **inner radius** of D as

$$r_D = \inf_{u \in \partial D'} H_0(u, \bar{u})^{1/2},$$

where D' is a bounded symmetric domain in standard form biholomorphically equivalent to D . By Theorem 2 of (4), if D is a Cartan classical domain, then

$$(7) \quad c_D = \frac{1}{r_D} = \begin{cases} \sqrt{2/(m+n)} & \text{if } D \in R_I, \\ \sqrt{2/(n+1)} & \text{if } D \in R_{II}, \\ \sqrt{1/(n-1)} & \text{if } D \in R_{III}, \\ \sqrt{2/n} & \text{if } D \in R_{IV}. \end{cases}$$

In particular, if D is the unit polydisk, then $c_D = 1$. If $D = \mathbb{B}^n$, then $c_D = \sqrt{2/(n+1)}$.

In (26), formula (7) was extended to include the case of the exceptional domains:

$$c_D = \frac{1}{r_D} = \begin{cases} 1/\sqrt{6} & \text{if } D = R_V, \\ 1/3 & \text{if } D = R_{VI}. \end{cases}$$

Furthermore, by Theorem 3 of (4) extended to include the exceptional domains, if $D = D_1 \times \cdots \times D_k$ is in standard form, then $c_D = \max_{1 \leq j \leq k} c_{D_j}$.

In the following proposition, for each of the four types of Cartan classical domains, we give explicit functions with range contained in $\mathbb{D}$ which are extremal in the sense that their semi-norms are equal to the Bergman constants of the domains. The proof of the following lemma can be found in (4) (Lemma 2 and Lemma 3).

Lemma 4.1. (4) (a) *Let Z be a matrix such that $I - ZZ^*$ is positive definite. Then each row of Z has norm less than 1.*

(b) *If $D \in R_{IV}$, then for each $z = (z_1, \dots, z_n) \in D$ and each distinct $r, s \in \{1, \dots, n\}$, $z_r \pm iz_s \in \mathbb{D}$.*

Proposition 4.1. (a) *If D is a Cartan classical domain in R_I , R_{II} , or R_{III} , then canonical projections $p_j(z) = z_j$ (for $j = 1, \dots, \dim(D)$) map D into $\mathbb{D}$ and have Bloch semi-norm equal to c_D .*

(b) *If $D \in R_{IV}$, then for each pair of distinct indices $r, s \in \{1, \dots, n\}$, for $j = 1, 2$, the functions $p_{r,s}^j$ defined on D by $p_{r,s}^1(z) = z_r + iz_s$ and $p_{r,s}^2(z) = z_r - iz_s$ have range contained in $\mathbb{D}$ and semi-norm equal to c_D .*

For easier reference, we shall refer to the functions in (b) as **modified projection maps**.

Proof. (a) The projections p_j map D into $\mathbb{D}$ by part (a) of Lemma 4.1. Represent the elements of $D \in R_I \cup R_{II} \cup R_{III}$ as vectors by considering the free variables in their matrix representation arranged rowwise. If $Z \in D$ with $D \in R_I$, then each entry of Z is a free variable and thus $Z = (z_{1,1}, z_{1,2}, \dots, z_{1,n}, z_{2,1}, z_{2,2}, \dots, z_{m,n})$. If $j \in \{1, \dots, mn\}$, let q and r be the unique integers such that $j = nq + r$ with $0 \leq q < m$ and $0 \leq r < n$ or $q = m$ and $r = 0$. Define e_j to be the unit vector whose j th coordinate is 1 and all others are 0, which corresponds to the matrix $E_{q,r}$ whose (q,r) -entry is 1 and all other entries are 0. Then

$$H_0(e_j, \overline{e_j}) = \left(\frac{m+n}{2} \right) \text{Trace}(E_{q,r} E_{q,r}^*) = \frac{m+n}{2}.$$

Furthermore, $|\nabla p_j(0)e_j| = 1$ so that $Q_{p_j}(0) = \sqrt{\frac{2}{m+n}} = c_D$.

If $Z \in D$ with $D \in R_{II}$, then the free variables are the entries on and above the main diagonal. Thus, for $j \in \{1, \dots, n(n+1)/2\}$ and the appropriate choice of indices h and k corresponding to j , we have

$$H_0(e_j, \overline{e_j}) = \left(\frac{n+1}{2} \right) \text{Trace}(E_{h,k} E_{h,k}^*) = \frac{n+1}{2}.$$

Furthermore, $|\nabla p_j(0)e_j| = 1$ so that $Q_{p_j}(0) = \sqrt{\frac{2}{n+1}} = c_D$.

If $Z \in D$ with $D \in R_{III}$, then the free variables are the entries above the main diagonal. For $j \in \{1, \dots, n(n-1)/2\}$ and the appropriate choice of indices h and k corresponding to j , let $X_{h,k} = E_{h,k} - E_{k,h} \in \partial D$. Then

$$H_0(e_j, \overline{e_j}) = \left(\frac{n-1}{2} \right) \text{Trace}(X_{h,k} X_{h,k}^*) = n-1.$$

Furthermore, $|\nabla p_j(0)e_j| = 1$ so that $Q_{p_j}(0) = \frac{1}{\sqrt{n-1}} = c_D$.

(b) Let $D \in R_{IV}$. The range of the modified projection maps $p_{r,s}^1$ and $p_{r,s}^2$ is contained in $\mathbb{D}$ by part (b) of Lemma 4.1. For $j = 1, 2$, let $e_{r,s}^j$ be the vector in $\mathbb{C}^n$ that has $1/2$ in the r -th place, $(-1)^j i/2$ in the s -th place, and 0 elsewhere. Thus $e_{r,s}^j \in \partial D$ and $H_0(e_{r,s}^j, \overline{e_{r,s}^j}) = n/2$. Moreover, $\nabla p_{r,s}^j(0)$ is the vector that has 1 in the r -th place, $(-1)^{j+1} i$ in the s -th place, and 0 elsewhere, so $|\nabla p_{r,s}^j(0)e_{r,s}^j| = 1$ and thus $Q_{p_{r,s}^j}(0) = \sqrt{\frac{2}{n}} = c_D$, completing the proof. $\square$

Corollary 4.1. If $D = D_1 \times \dots \times D_k$ is a bounded symmetric domain, with $D_1, \dots, D_k$ Cartan classical domains, then the projection maps or the modified projection maps for the case of a factor in R_{IV} of D onto $\mathbb{D}$ have semi-norms $c_{D_1}, \dots, c_{D_k}$ in this order, repeated as many times as the dimensions of the factors $D_1, \dots, D_k$.

Proof. Since the Bergman metric of $D_1 \times \cdots \times D_k$ is the direct sum of the Bergman metrics of the individual factors, the result follows at once from Proposition 4.1. $\square$

In Section 6, we shall prove that if $D = D_1 \times \cdots \times D_k$, where each D_j is a Cartan classical domain, and C_φ is an isometry on $\mathcal{B}$, then the components $\varphi_j = p_j \circ \varphi$ or the functions $\varphi_r + i\varphi_s$ and $\varphi_r - i\varphi_s$ (which we call **modified components**) must have Bloch semi-norms equal to $c_{D_1}, \dots, c_{D_k}$, respectively, repeated according to the dimension of the factors.

5. THE HOLOMORPHIC SELF MAPS ON A BOUNDED HOMOGENEOUS DOMAIN

A bounded homogeneous domain D considered as a Riemannian manifold under the structure induced by the Bergman metric is a length space. Therefore, given $\varphi \in H(D)$ viewed as a map between Riemannian manifolds, we may apply Property 1.8 bis of (13). We obtain the following interpretation of the Bergman constant.

Proposition 5.1. *Any function $\varphi \in H(D)$ is Lipschitz as a mapping from (D, ρ) into itself and the Bergman constant B_φ is the Lipschitz number of φ . That is*

$$B_\varphi = \sup_{z \neq w} \frac{\rho(\varphi(z), \varphi(w))}{\rho(z, w)}.$$

Using Proposition 5.1 and the invariance of the Bergman distance under the action of an automorphism of D , we obtain

Corollary 5.1. Let D be a bounded homogeneous domain and let $\varphi \in H(D)$. Then for all $S \in \text{Aut}(D)$, $B_{\varphi \circ S} = B_\varphi = B_{S \circ \varphi}$.

The proof of the following result is analogous to that of Theorem 3.3 where the Euclidean distance is replaced by the Bergman distance.

Corollary 5.2. Let D be a bounded homogeneous domain. If $\{\varphi_n\}$ is a sequence in $H(D)$ converging to a function $\varphi \in H(D)$ uniformly on compact subsets of D , then $B_\varphi \leq \liminf_{n \rightarrow \infty} B_{\varphi_n}$.

Theorem 5.1. *Let D be a bounded homogeneous domain and let $\varphi \in H(D)$ be such that $\varphi(0) = 0$ and suppose there exists a sequence $\{S_j\}$ in $\text{Aut}(D)$ such that $\{\varphi \circ S_j\}$ converges locally uniformly in D to the identity. If the Bergman constant of φ does not exceed 1, then C_φ is an isometry on $\mathcal{B}$.*

Proof. First observe that, by Corollary 5.2 and Corollary 5.1, $1 \leq B_{\varphi \circ S_j} = B_\varphi$. Thus, from the hypothesis it follows that $B_\varphi = 1$. If $f \in \mathcal{B}$, then $f \circ \varphi \circ S_n \rightarrow f$ locally uniformly in D , so by Theorem 3.3 and the invariance of the Bloch semi-norm under composition of automorphisms, we obtain

$$\beta_f \leq \liminf_{n \rightarrow \infty} \beta_{f \circ \varphi \circ S_n} = \beta_{f \circ \varphi} \leq B_\varphi \beta_f = \beta_f.$$

Thus, since $\varphi(0) = 0$, $\|f \circ \varphi\|_{\mathcal{B}} = |f(0)| + \beta_{f \circ \varphi} = |f(0)| + \beta_f = \|f\|_{\mathcal{B}}$. $\square$

In (17), Korányi proved the following version of the Schwarz lemma for bounded symmetric domains.

Theorem 5.2. (17) *Let D be a bounded symmetric domain of rank ℓ and let $\varphi \in H(D)$. Then for all $z, w \in D$, $\rho(\varphi(z), \varphi(w)) \leq \ell^{1/2} \rho(z, w)$, where ρ is the Bergman distance on D . The constant $\ell^{1/2}$ is the best possible.*

The rank of D is roughly the dimension of the largest polydisk that can be embedded in the tangent space at each point of D , considered as a complex manifold under the Bergman metric. In particular, if $D = \mathbb{D}^n$, then $\ell = n$ so that $\max_{\varphi \in H(D)} B_\varphi = \sqrt{n}$. If D is the unit ball, $\ell = 1$ so that $\max_{\varphi \in H(D)} B_\varphi = 1$. Thus, in the statement of Theorem 5.1 for the case $D = \mathbb{B}_n$, the hypothesis on B_φ is unnecessary.

Theorem 5.1 allows us to obtain non-trivial examples of isometric composition operators if D is a bounded homogeneous domain having $\mathbb{D}$ as a factor.

Example 5.1. Let $D = D_1 \times \mathbb{D}$, where D_1 is a bounded homogeneous domain containing 0. Let $\phi \in H(\mathbb{D})$ be such that $\phi(0) = 0$, $\beta_\phi = 1$, and suppose ϕ is not a rotation. Let $U \in \text{Aut}(D_1)$ such that $U(0) = 0$. For $z \in D_1$ and $\zeta \in \mathbb{D}$, define $\varphi(z, \zeta) = (U(z), \phi(\zeta)) \in D$. By the equivalence of (a) and (g) in Theorem 1.1, there exists a sequence $\{T_j\}$ in $\text{Aut}(\mathbb{D})$ such that $\phi \circ T_j(\zeta) \rightarrow \zeta$ locally uniformly in $\mathbb{D}$. Then the sequence $\{S_j\}$ defined by $S_j(z, \zeta) = (U^{-1}(z), T_j(\zeta))$ is in $\text{Aut}(D)$ and $\{\varphi \circ S_j\}$ converges locally uniformly to the identity in D . Since the Bergman constants of U and ϕ are both 1, then so is the Bergman constant of φ . By Theorem 5.1, C_φ is an isometry on $\mathcal{B}$.

6. SOME NECESSARY CONDITIONS

In this section, we derive some necessary conditions for $\varphi \in H(D)$ to be the symbol of an isometric composition operator on the Bloch space when D is a bounded symmetric domain in standard form with no exceptional factors. These conditions are analogous to those found in (6) in the special case of the unit polydisk for domains that do not have factors in R_{IV} . A modification is needed when dealing with domains that contain factors in R_{IV} . The key tool in the proof of Theorem 6.1 below is the existence of an automorphism interchanging any prescribed point of D with 0.

Lemma 6.1. *Let D be a bounded symmetric domain in standard form. For each $a \in D$, there exists an involution $\psi_a \in \text{Aut}(D)$ such that $\psi_a(a) = 0$.*

The proof is an immediate consequence of the results in (15), pp. 170, 301, 311.

Theorem 6.1. *Let $D = D_1 \times \cdots \times D_k$ be a bounded symmetric domain in standard form. Let $\varphi = (\varphi_1, \dots, \varphi_n) \in H(D)$ such that C_φ is an isometry on $\mathcal{B}$. Then:*

- (a) *The components $\varphi_1, \dots, \varphi_n$ are linearly independent.*
- (b) *If D does not contain any exceptional factors, then $\varphi(0) = 0$.*
- (c) *If none of the factors of D is in R_{IV} , then the components of φ have semi-norms equal to $c_{D_1}, \dots, c_{D_k}$, repeated according to the dimension of each factor.*
- (d) *If D has factors in R_{IV} , then for each $D_\ell \in R_{IV}$ and each pair r, s of distinct indices, with $\sum_{i=1}^{\ell-1} \dim(D_i) < r, s \leq \sum_{i=1}^{\ell} \dim(D_i)$, the modified components $\varphi_r + i\varphi_s, \varphi_r - i\varphi_s \in H(D, \mathbb{D})$ and have semi-norm equal to c_{D_ℓ} .*

Proof. To prove (a), assume there exist $k = 1, \dots, n$ and $\alpha_j \in \mathbb{C}$ for $j \neq k$ such that $\varphi_k = \sum_{j \neq k} \alpha_j \varphi_j$. Then the function f defined by $f(z) = z_k - \sum_{j \neq k} \alpha_j z_j$ is bounded on D (and hence a Bloch function) and has nonzero Bloch norm, but $f \circ \varphi$ is identically zero. Thus C_φ is not an isometry on $\mathcal{B}$.

To prove (b), assume each of the factors of D is a Cartan classical domain. Set $\varphi(0) = a$ and write $a = (a_1, \dots, a_n)$, with each $a_j \in \mathbb{C}$. Fix $j = 1, \dots, n$. By Lemma 6.1, we may consider $\psi_j(z) = p_j \circ \psi_a$, where p_j is the projection map onto the

j -th coordinate and ψ_a is an automorphism of D mapping a to 0 and 0 to a . Then, by the invariance of the semi-norm under composition of automorphisms of D and by Corollary 4.1, if $D_\ell \notin R_{IV}$ and $\dim(D_1 \times \cdots \times D_{\ell-1}) < j \leq \dim(D_1 \times \cdots \times D_\ell)$, then $\beta_{\psi_j} = \beta_{p_j} = c_{D_\ell}$. On the other hand, $\beta_{\psi_j \circ \varphi} \leq c_{D_\ell}$. Since $\|\psi_j \circ \varphi\|_{\mathcal{B}} = \|\psi_j\|_{\mathcal{B}}$, and $\psi_j(a) = 0$, we deduce $|a_j| + c_{D_\ell} = \beta_{\psi_j \circ \varphi} \leq c_{D_\ell}$. Hence $a_j = 0$ and $\beta_{\psi_j \circ \varphi} = c_{D_\ell}$.

Next assume $D_\ell \in R_{IV}$ and $\dim(D_1 \times \cdots \times D_{\ell-1}) < r, s \leq \dim(D_1 \times \cdots \times D_\ell)$, with $r \neq s$. Define $\psi_{r,s}^h = p_{r,s}^h \circ \psi_a$, where $p_{r,s}^h$ ($h = 1, 2$) are the modified projection maps defined in Proposition 4.1. Then $\beta_{\psi_{r,s}^h \circ \varphi} \leq c_{D_\ell}$, and since $\|\psi_{r,s}^h \circ \varphi\|_{\mathcal{B}} = \|\psi_{r,s}^h\|_{\mathcal{B}}$, and $\psi_{r,s}^h(a) = 0$, by Corollary 4.1 we deduce

$$|a_r + (-1)^{h+1}ia_s| + c_{D_\ell} = \beta_{\psi_{r,s}^h \circ \varphi} \leq c_{D_\ell}.$$

Hence $a_r + (-1)^{h+1}ia_s = 0$ for $h = 1, 2$, which implies that $a_r = a_s = 0$. Consequently, $\varphi(0) = 0$.

Moreover, $\beta_{\varphi_j} = \beta_{p_j \circ \varphi} = \beta_{p_j} = c_{D_\ell}$ if $D_\ell \notin R_{IV}$, while $\beta_{\varphi_r + (-1)^{h+1}i\varphi_s} = \beta_{p_{r,s}^h \circ \varphi} = \beta_{p_{r,s}^h} = c_{D_\ell}$ if $D_\ell \in R_{IV}$, proving (c) and (d). $\square$

Remark 6.1. We suspect that if D is any bounded symmetric domain in standard form, then the condition $\varphi(0) = 0$ holds, but we have not been able to verify it in the case an irreducible factor is an exceptional domain because our proof relies on our ability to find explicit functions mapping into the unit disk that are extremal with respect to the Bloch semi-norm and attain the semi-norm at 0 (these were the projection maps for domains in R_I, R_{II} , and R_{III} , and the modified projection maps for domains in R_{IV}). We have not been able to determine any functions defined on the exceptional domains that have these properties.

Thus, we propose the following

Conjecture. Let $\varphi \in H(D)$, where $D = D_1 \times \cdots \times D_k$ is a bounded symmetric domain in standard form. Then C_φ is an isometry on $\mathcal{B}$ if and only if $\varphi(0) = 0$, $B_\varphi = 1$, and the components or the modified components of φ have semi-norm equal to $c_{D_1}, \dots, c_{D_k}$, repeated according to the dimension of each factor.

7. THE SPECTRUM OF C_φ IN THE CASE OF THE POLYDISK

In this section, we study the spectrum of a large class of isometric composition operators on the Bloch space of the unit polydisk $\mathbb{D}^n$. As a consequence, we obtain a complete classification of the spectrum in the case of the unit disk.

First, we recall pertinent facts of the spectrum of bounded linear operators on a complex Banach space. These results can all be found in a standard text of functional analysis such as (10).

A bounded linear operator A on a complex Banach space E is invertible if and only if it has dense range and is *bounded below*, that is, there exists $c > 0$ such that for all $x \in E$, $\|Ax\| \geq c\|x\|$. By the Inverse Mapping Theorem, for invertibility it is sufficient for A to be bijective.

The **resolvent** of a bounded linear operator A on a complex Banach space E is defined as

$$\rho(A) = \{\lambda \in \mathbb{C} : A - \lambda I \text{ is invertible}\},$$

where I is the identity operator. The **spectrum** of A is defined as $\sigma(A) = \mathbb{C} \setminus \rho(A)$. The **approximate point spectrum**, a subset of the spectrum, is defined as

$$\sigma_{ap}(A) = \{\lambda \in \mathbb{C} : A - \lambda I \text{ is not bounded below}\}.$$

The spectrum of an operator A is a non-empty, compact subset of $\mathbb{C}$ contained in the closed disk centered at the origin of radius equal to the operator norm of A . In particular, the spectrum of an isometry is contained in $\overline{\mathbb{D}}$, since the operator norm is 1. Furthermore $\partial\sigma(A) \subseteq \sigma_{ap}(A)$ (see (10), Proposition 6.7).

To prove our results on the spectrum, we need the following lemma.

Lemma 7.1. *Let E be a complex Banach space and suppose $T : E \rightarrow E$ is an isometry. If T is invertible, then $\sigma(T) \subseteq \partial\mathbb{D}$. If T is not invertible, then $\sigma(T) = \overline{\mathbb{D}}$.*

Proof. Suppose T is an invertible isometry on E . Then $0 \notin \sigma(T)$, and so the function $z \mapsto z^{-1}$ is analytic in some neighborhood of $\sigma(T)$. By the Spectral Mapping Theorem (see (10), p. 204), we have $\sigma(f \circ T) = f(\sigma(T))$, and so

$$\sigma(T^{-1}) = \sigma(T)^{-1} = \{\lambda^{-1} : \lambda \in \sigma(T)\}.$$

Since T^{-1} exists and is an isometry, we have $\sigma(T^{-1}) \subseteq \overline{\mathbb{D}}$. Therefore $\sigma(T) \subseteq \partial\mathbb{D}$.

Next, suppose T is not invertible. In order to prove that $\sigma(T) = \overline{\mathbb{D}}$, it suffices to show that $\overline{\mathbb{D}} \subseteq \sigma(T)$. For $\lambda \in \mathbb{D}$, $T - \lambda I$ is bounded below by $1 - |\lambda|$. Thus, $\lambda \notin \sigma_{ap}(T)$. We deduce that $\partial\sigma(T) \subseteq \sigma_{ap}(T) \subseteq \partial\mathbb{D}$.

Since T is not invertible, $0 \in \sigma(T)$. Assume $\lambda \in \overline{\mathbb{D}} \cap \rho(T)$. Note that $\lambda \notin \partial\sigma(T)$ since $\partial\sigma(T) = \sigma(T) \cap \overline{\rho(T)}$. Consider $\Gamma = \{t\lambda : t \in [0, \infty)\}$, the radial line through λ . Since $\sigma(T)$ is closed, there exists $t \in [0, 1)$ such that $t\lambda \in \partial\sigma(T)$. This contradicts the fact that $\partial\sigma(T) \subseteq \partial\mathbb{D}$. Consequently, $\overline{\mathbb{D}} \cap \rho(T) = \emptyset$, whence $\overline{\mathbb{D}} \subseteq \sigma(T)$. $\square$

We use Lemma 7.1 to obtain a classification of the spectrum for a large class of symbols of isometric composition operators for the polydisk. We first describe the spectrum of isometric composition operators with surjective, non-automorphic symbol.

Proposition 7.1. *If φ is the symbol of an isometric composition operator such that $\varphi \notin \text{Aut}(\mathbb{D}^n)$ and φ is onto, then $\sigma(C_\varphi) = \overline{\mathbb{D}}$.*

Proof. By Lemma 7.1, it suffices to show that $0 \in \sigma(C_\varphi)$. Since C_φ is an isometry, it is necessarily injective. Thus, we must prove that C_φ is not surjective. Arguing by contradiction, suppose C_φ is surjective. Since φ is onto but not an automorphism, φ is not 1-1. Thus there exist distinct $\zeta, \eta \in \mathbb{D}^n$ such that $\varphi(\zeta) = \varphi(\eta)$. In particular, there exists $k = 1, \dots, n$ such that $\zeta_k \neq \eta_k$.

Let $p_k : \mathbb{D}^n \rightarrow \mathbb{D}$ be the projection map onto the k^{th} coordinate, and define the Bloch function $g(z) = p_k(z) - \eta_k$. Note that g is non-vanishing at ζ . Since C_φ is surjective, there exists a Bloch function f such that $f \circ \varphi = g$. In particular $f(\varphi(\zeta)) = g(\zeta) \neq 0$. On the other hand, $f(\varphi(\zeta)) = f(\varphi(\eta)) = g(\eta) = 0$, a contradiction. Thus, C_φ is not surjective. Therefore $\sigma(C_\varphi) = \overline{\mathbb{D}}$. $\square$

Observe that the class of symbols described in Proposition 7.1 includes those of the form $\varphi = (\varphi_1, \dots, \varphi_n)$ with each φ_k dependent on a distinct single complex variable as described in Theorem 1.1(f).

Next, we consider the isometric composition operators induced by a particular type of automorphic symbol. Let $\varphi(z) = (\lambda_1 z_1, \dots, \lambda_n z_n)$, with $|\lambda_j| = 1$ for $j = 1, \dots, n$. We define $\text{ord}(\lambda_j)$ to be the smallest nonnegative integer k such that $\lambda_j^k = 1$. We have the following classification of the spectrum of C_φ for this class of automorphisms.

Proposition 7.2. *Let $\varphi(z) = (\lambda_1 z_1, \dots, \lambda_n z_n)$ such that $|\lambda_j| = 1$ for $j = 1, \dots, n$. Then*

$$\sigma(C_\varphi) = \begin{cases} \partial\mathbb{D}, & \text{if } \text{ord}(\lambda_j) = \infty \text{ for some } j, \\ G, & \text{if } \text{ord}(\lambda_j) < \infty \text{ for all } j, \end{cases}$$

where G is the finite cyclic group generated by $\lambda_1, \dots, \lambda_n$.

Proof. The induced composition operator C_φ is clearly invertible since $\varphi^{-1}(z) = (\lambda_1^{-1} z_1, \dots, \lambda_n^{-1} z_n)$ and $C_\varphi^{-1} = C_{\varphi^{-1}}$. So, by Lemma 7.1, we have $\sigma(C_\varphi) \subseteq \partial\mathbb{D}$.

Observe that the group G generated by $\lambda_1, \dots, \lambda_n$ is

$$\{\lambda_1^{k_1} \cdots \lambda_n^{k_n} : k_1, \dots, k_n \in \mathbb{N} \cup \{0\}\} \subseteq \partial\mathbb{D}.$$

For $k_1, \dots, k_n \in \mathbb{N} \cup \{0\}$, the function $f(z) = z_1^{k_1} \cdots z_n^{k_n}$ is Bloch and $(f \circ \varphi)(z) = f(\lambda_1 z_1, \dots, \lambda_n z_n) = \lambda_1^{k_1} \cdots \lambda_n^{k_n} f(z)$. Thus, f is an eigenfunction of C_φ with eigenvalue $\lambda_1^{k_1} \cdots \lambda_n^{k_n}$. Thus, $\overline{G} \subseteq \sigma(C_\varphi)$.

If there exists $j = 1, \dots, n$ such that $\text{ord}(\lambda_j) = \infty$, then G is a dense subset of $\partial\mathbb{D}$. Thus, $\partial\mathbb{D} = \overline{G} \subseteq \sigma(C_\varphi)$, and hence $\sigma(C_\varphi) = \partial\mathbb{D}$. If $\text{ord}(\lambda_j) < \infty$ for all $j = 1, \dots, n$, then G is a finite cyclic group of order equal to the least common multiple of the orders of $\lambda_1, \dots, \lambda_n$. We wish to show that $\sigma(C_\varphi) \subseteq G$. Let $\mu \in \partial\mathbb{D} \setminus G$. The invertibility of $C_\varphi - \mu I$ is equivalent to showing that for all Bloch functions g , there exists a unique Bloch function f such that $C_\varphi(f) - \mu f = g$.

If α is the order of G , then $\varphi^\alpha(z) := \underbrace{(\varphi \circ \cdots \circ \varphi)}_{\alpha\text{-times}}(z) = z$ for all $z \in \mathbb{D}^n$, so

$C_\varphi - \mu I$ is invertible if and only if for every Bloch function g the following system has a unique solution f in the Bloch space:

$$(8) \quad \begin{array}{rcl} f(\varphi(z)) & - & \mu f(z) = g(z) \\ f(\varphi^2(z)) & - & \mu f(\varphi(z)) = g(\varphi(z)) \\ & \vdots & \\ f(z) & - & \mu f(\varphi^{\alpha-1}(z)) = g(\varphi^{\alpha-1}(z)). \end{array}$$

Equivalently, (8) can be posed as the matrix equation $Ax = b$ where

$$A = \begin{bmatrix} -\mu & 1 & 0 & 0 & \cdots & 0 \\ 0 & -\mu & 1 & 0 & \cdots & 0 \\ \vdots & 0 & \ddots & \ddots & & \vdots \\ \vdots & & \ddots & \ddots & \ddots & \vdots \\ 0 & & & \ddots & \ddots & 1 \\ 1 & 0 & \cdots & \cdots & 0 & -\mu \end{bmatrix}, x = \begin{bmatrix} f(z) \\ f(\varphi(z)) \\ \vdots \\ f(\varphi^{\alpha-2}(z)) \\ f(\varphi^{\alpha-1}(z)) \end{bmatrix}, \text{ and } b = \begin{bmatrix} g(z) \\ g(\varphi(z)) \\ \vdots \\ g(\varphi^{\alpha-2}(z)) \\ g(\varphi^{\alpha-1}(z)) \end{bmatrix}.$$

Direct calculation shows that $\det(A) = (-1)^\alpha(\mu^\alpha - 1) \neq 0$, since $\mu \notin G$. Thus, $C_\varphi - \mu I$ is invertible. For $\mu \notin G$, the unique solution f of (8) is a (finite) linear combination of the functions $g \circ \varphi^{j-1}$, for $j = 1, \dots, \alpha$, each of which is Bloch. Thus f is Bloch. Therefore $\sigma(C_\varphi) = G$. $\square$

We now study the spectrum in the case when φ is an automorphism that permutes the coordinates.

Proposition 7.3. *Let $\varphi(z) = (z_{\tau(1)}, \dots, z_{\tau(n)})$, where $\tau \in S_n$ is decomposed into a product of disjoint cycles $c_1, \dots, c_\ell$ of order $\alpha_1, \dots, \alpha_\ell$, respectively.*

- (a) If λ is an α_j -th root of unity for some $j = 1, \dots, \ell$, then λ is an eigenvalue of C_φ .
- (b) The spectrum of C_φ is the group of the α -th roots of unity, where α is the order of τ .

Proof. To prove (a), assume λ is an α_j -th root of unity for some $j = 1, \dots, \ell$. To prove that λ is an eigenvalue we need to show that there exists a nonzero $f \in \mathcal{B}$ such that

$$(9) \quad f(\varphi(z)) = \lambda f(z) \text{ for all } z \in \mathbb{D}^n.$$

We are going to show that, in fact, there is a linear function f satisfying this condition. Let $f(z) = \sum_{j=1}^n x_j z_j$, where the coefficients x_j are to be determined. Equation (9) is equivalent to a matrix equation $Bx = 0$, where x is the column vector with coordinates $x_1, \dots, x_n$ and B is a matrix with all diagonal entries $-\lambda$ and whose rows (and columns) contain an entry 1 and all other entries 0. The matrix B can be permuted to yield the block diagonal matrix

$$A = \begin{bmatrix} A_1 & O & \dots & O \\ O & A_2 & \dots & O \\ \vdots & & \ddots & \vdots \\ O & O & \dots & A_n \end{bmatrix}$$

where each matrix A_k has order $\alpha_k \times \alpha_k$ and contains all diagonal entries 1, and each row contains (besides the entry 1) one entry $-\lambda$ and all other entries 0. The matrix A_k corresponds to the cycle c_k . It is easy to see that $\det(A_j) = 1 - \lambda^{\alpha_j} = 0$, so that $\det(A) = \prod_{k=1}^\ell \det(A_k) = 0$. Thus B itself is singular, and so $Bx = 0$ has nontrivial solutions.

To prove (b), assume μ is not an α -th root of unity. Arguing as in the proof of Proposition 7.2, given a function $g \in \mathcal{B}$, we form the system (8), whose coefficient matrix has nonzero determinant. Thus, for any function $g \in \mathcal{B}$ there exists a unique $f \in \mathcal{B}$ such that $f \circ \varphi - \mu f = g$.

On the other hand, if μ is an α -th root of unity, then by row reduction it is easy to see that the solvability of the system (8) reduces to the condition

$$\begin{aligned} &g(\varphi^{\alpha-2}(z)) + \mu g(\varphi^{\alpha-3}(z)) + \mu^2 g(\varphi^{\alpha-4}(z)) + \dots \\ &\dots + \mu^{\alpha-3} g(\varphi(z)) + \mu^{\alpha-2} g(z) + \mu^{\alpha-1} g(\varphi^{\alpha-1}(z)) \equiv 0. \end{aligned}$$

By choosing g to be a polynomial of sufficiently large degree, it is possible to find one for which identity (7) fails. Thus, $\mu \in \sigma(C_\varphi)$. $\square$

As observed in Section 2, the automorphisms of the polydisk that fix the origin have the representation $\varphi(z) = (\lambda_1 z_{\tau(1)}, \dots, \lambda_n z_{\tau(n)})$ where $|\lambda_j| = 1$ for all $j \in \{1, \dots, n\}$, and $\tau \in S_n$. The arguments used in the proofs of Proposition 7.2 and Proposition 7.3 carry over to this general case, where α is taken to be the least common multiple of the orders of $\lambda_1, \dots, \lambda_n$, and τ . We deduce the following result:

Theorem 7.1. *Let φ be the symbol of an isometric composition operator on the Bloch space for the polydisk.*

- (a) If $\varphi \notin \text{Aut}(\mathbb{D}^n)$ and φ is onto, then $\sigma(C_\varphi) = \overline{\mathbb{D}}$.

- (b) If $\varphi \in \text{Aut}(\mathbb{D}^n)$, let $\lambda_1, \dots, \lambda_n \in \partial\mathbb{D}$ and let $\tau \in S_n$ be such that $\varphi(z) = (\lambda_1 z_{\tau(1)}, \dots, \lambda_n z_{\tau(n)})$.
 (i) If at least one of the λ_j has infinite order, then $\sigma(C_\varphi) = \partial\mathbb{D}$.
 (ii) If each λ_j has finite order, then $\sigma(C_\varphi)$ is the cyclic group Γ generated by $\lambda_1, \dots, \lambda_n$ and by the m^{th} roots of unity, where m is the order of τ . Furthermore, each element of the group G generated by $\lambda_1, \dots, \lambda_n$ is an eigenvalue.

We are not aware of any non-onto symbols of isometric composition operators on $\mathbb{D}^n$. Thus, we propose the following

Conjecture. Let $\varphi \in H(\mathbb{D}^n)$ be the symbol of an isometric composition operator on $\mathcal{B}$.

- (a) If $\varphi \notin \text{Aut}(\mathbb{D}^n)$, then $\sigma(C_\varphi) = \overline{\mathbb{D}}$.
 (b) If $\varphi \in \text{Aut}(\mathbb{D}^n)$, so that $\varphi(z) = (\lambda_1 z_{\tau(1)}, \dots, \lambda_n z_{\tau(n)})$ where $|\lambda_j| = 1$ for $j = 1, \dots, n$ and $\tau \in S_n$, then

$$\sigma(C_\varphi) = \begin{cases} \partial\mathbb{D}, & \text{if } \text{ord}(\lambda_j) = \infty \text{ for some } j, \\ \Gamma, & \text{if } \text{ord}(\lambda_j) < \infty \text{ for all } j, \end{cases}$$

where Γ is the finite cyclic group generated by $\lambda_1, \dots, \lambda_n$ and by the m^{th} roots of unity, with $m = \text{ord}(\tau)$.

From Theorem 7.1 and Theorem 1.1, which gives a complete characterization of the symbols of the isometric composition operators on the Bloch space of the unit disk, we deduce a complete classification of the spectrum in the one-dimensional case.

Theorem 7.2. Let φ be the symbol of an isometric composition operator on the Bloch space of the unit disk.

- (a) If φ is not a rotation, then $\sigma(C_\varphi) = \overline{\mathbb{D}}$.
 (b) If $\varphi(z) = \lambda z$ with $|\lambda| = 1$, then

$$\sigma(C_\varphi) = \begin{cases} \partial\mathbb{D}, & \text{if } \text{ord}(\lambda) = \infty, \\ G, & \text{if } \text{ord}(\lambda) < \infty, \end{cases}$$

where G is the cyclic group generated by λ .

Proof. The proof is the direct application of Theorem 7.1 to the case of the unit disk, since in this case the symbols of the isometric composition operators are onto (see part (c) of Theorem 1.1). $\square$

8. OPEN QUESTIONS

We conclude the paper with the following open questions.

- (1) In the case when D is the unit disk or the unit ball, it is well known (see (27) (Theorem 5.1.7) and (28) (Theorem 3.9)) that for any $z, w \in D$,

$$\rho(z, w) = \sup\{|f(z) - f(w)| : f \in \mathcal{B}, \beta_f \leq 1\}.$$

On the other hand, by the characterization of the semi-norm as a Lipschitz number, it follows immediately that for any bounded homogeneous domain D and $z, w \in D$,

$$\sup\{|f(z) - f(w)| : f \in \mathcal{B}, \beta_f \leq 1\} \leq \rho(z, w).$$

For which domains D does the opposite inequality hold?

If D is such a domain and $\varphi \in H(D)$, we obtain a sharper lower estimate of the norm of C_φ . Indeed, using a normal family argument, we can find $f \in \mathcal{B}$ such that $\beta_f = 1$ and $\rho(\varphi(0), 0) = |f(\varphi(0)) - f(0)|$. Letting $g(z) = f(z) - f(0)$, we obtain a Bloch function g fixing 0 with norm 1 and such that $\rho(\varphi(0), 0) = |g(\varphi(0))|$. Thus $\|C_\varphi(g)\|_{\mathcal{B}} \geq |g(\varphi(0))| = \rho(\varphi(0), 0)$. Since as observed in Theorem 3.2, $\|C_\varphi\| \geq 1$, we obtain $\|C_\varphi\| \geq \max\{1, \rho(\varphi(0), 0)\}$. In the case of the unit ball this is the lower estimate in Corollary 3.1.

At the end of Section 5, we saw that if D is a bounded homogeneous domain having D as a factor, then it is possible to construct non-trivial examples of symbols of isometric composition operators.

(2) Are there any non-trivial examples of isometric composition operators on the Bloch space for domains D which do not contain the disk as a factor? In particular, are there any symbols of isometric composition operators on the Bloch space for the unit ball that are not unitary transformations?

In the case of the polydisk, although there is a large class of non-trivial isometries C_φ on $\mathcal{B}$, we haven't been able to find any symbols φ such that $B_\varphi > 1$.

(3) Are there any symbols φ for which C_φ is an isometry on $\mathcal{B}$ and $B_\varphi > 1$ if D is not the unit ball?

(4) If φ is a holomorphic self-map of a bounded homogeneous domain D in $\mathbb{C}^n$ with $n > 1$ and C_φ is an isometry on the Bloch space of D , does there exist a sequence $\{S_k\}$ in $\text{Aut}(D)$ such that $\{\varphi \circ S_k\}$ converges locally uniformly to the identity in D ?

The answer in (4) is affirmative if $D = \mathbb{D}$ by Theorem 1.1. The following is a weaker form of a positive answer in question (4) for the case of the polydisk (see Theorem 8 in (6)).

Theorem 8.1. *Let $\varphi \in H(\mathbb{D}^n)$ be such that C_φ is an isometry on $\mathcal{B}$. Then there exist sequences $\{T_m^j\}_{m \in \mathbb{N}}$ ($j = 1, \dots, n$) of automorphisms of $\mathbb{D}^n$ such that $\{(\varphi_1 \circ T_m^1, \dots, \varphi_n \circ T_m^n)\}$ converges uniformly on compact subsets to the identity in $\mathbb{D}^n$.*

In (6), the following simple example was given to show that the converse to Theorem 8.1 is false. Let $\varphi(z_1, z_2) = (z_1, z_1)$ for $|z_j| < 1$, $j = 1, 2$. Since the components of φ are equal, C_φ is not an isometry. On the other hand, $(\varphi_1(z_1, z_2), \varphi_2(T(z_1, z_2))) = (z_1, z_2)$, where $T(z_1, z_2) = (z_2, z_1)$.

In (6), the question on whether the components of the symbols φ of the isometric composition operators for the polydisk $\varphi \in H(\mathbb{D}^n)$ must depend on a single distinct complex variable was raised.

(5) Are there any other holomorphic maps φ which induce an isometric composition operator on the Bloch space of the polydisk besides those whose components are defined in terms of the isometries on the Bloch space on $\mathbb{D}$?

A negative answer to this question and Theorem 8.1 would yield a positive answer to question (4) for the case of the polydisk.

(6) Are there any isometric composition operators on the Bloch space of the polydisk whose symbols are not onto?

Definition 8.1. (11) A functional Banach space $\mathcal{Y}$ on a set X is called **algebraically consistent** if the only non-zero bounded linear functionals k on $\mathcal{Y}$ such that $k(fg) = k(f)k(g)$ whenever $f, g, fg \in \mathcal{Y}$ are point evaluations (i.e. functionals of the form $K_x(f) = f(x)$ for each $f \in \mathcal{Y}$, where $x \in X$).

In Theorem 1.6 of (11) it is shown that if $\mathcal{Y}$ is an algebraically consistent functional Banach space consisting of analytic functions on a set X whose interior is connected and dense in X and such that the elements of $\mathcal{Y}$ are analytic in the interior of X and continuous on X , and if φ is an analytic self-map of the interior of X such that C_φ is bounded and invertible on $\mathcal{Y}$, then φ is one-to-one and onto and $C_\varphi^{-1} = C_{\varphi^{-1}}$.

(7) Is the Bloch space of the polydisk algebraically consistent?

In the case of an affirmative answer, it would follow that if C_φ is an isometry on the Bloch space of the polydisk with $\varphi \notin \text{Aut}(\mathbb{D}^n)$, then C_φ is not invertible (so that $\sigma(C_\varphi) = \overline{\mathbb{D}}$), without assuming that φ is onto.

REFERENCES

- [1] J. M. Anderson, J. Clunie, Ch. Pommerenke, *On Bloch functions and normal functions*, J. Reine Angew. Math. **279** (1974), 12–37.
- [2] E. Cartan, *Sur les domaines bornés de l'espace de n variable complexes*, Abh. Math. Sem. Univ. Hamburg **11** (1935), 116–162.
- [3] J. Cima, *The basic properties of Bloch functions*, Internat. J. Math. & Math. Sci. (2) **3** (1979), 369–413.
- [4] J. M. Cohen, F. Colonna, *Bounded holomorphic functions on bounded symmetric domains*, Trans. Amer. Math. Soc. (1) **343** (1994), 135–156.
- [5] J. M. Cohen, F. Colonna, *Preimages of one-point sets of Bloch and normal functions*, Mediterr. J. Math. **3** (2006), 513–532.
- [6] J. M. Cohen, F. Colonna, *Isometric composition operators on the Bloch space in the polydisk*, Contemp. Math., **454** (2008), 9–21.
- [7] F. Colonna, *Characterisation of the isometric composition operators on the Bloch space*, Bull. Austral. Math. Soc., **72** (2005), 283–290.
- [8] F. Colonna, *Bloch and normal functions and their relation*, Rend. Circ. Mat. Palermo, Ser. II **XXXVIII** (1989), 161–180.
- [9] F. Colonna, *The Bloch constant of bounded analytic functions*, J. London Math. Soc. (2) **36** (1987), 95–101.
- [10] J. B. Conway, *A Course in Functional Analysis*, Springer-Verlag, New York, 1994.
- [11] C. Cowen, B. MacCluer, *Composition Operators on Spaces of Analytic Functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton (1995).
- [12] D. Drucker, *Exceptional Lie algebras and the structure of Hermitian symmetric spaces*, Mem. Amer. Math. Soc. **208** (1978), 1–207.
- [13] M. Gromov, *Structures Métriques pour les Variétés Riemanniennes*, Cedric/Fernand Nathan, Paris, 1981.
- [14] K. T. Hahn, *Holomorphic mappings of the hyperbolic space into the complex Euclidean space and the Bloch theorem*, Canad. J. Math. **27** (1975), 446–458.
- [15] S. Helgason, *Differential Geometry and Symmetric Spaces*, Academic Press, New York-London, 1962.
- [16] S. Kobayashi, *Hyperbolic manifolds and holomorphic mappings, an Introduction* (2nd ed.), World Scientific, London, 2005.
- [17] A. Korányi, *A Schwarz lemma for bounded symmetric domains*, Proc. Amer. Math. Soc., **17** (1966), 210–213.
- [18] S. G. Krantz, *Function Theory of Several Complex Variables*, John Wiley & Sons, New York, 1982.

- [19] S. G. Krantz, D. Ma, *Bloch functions on strongly pseudoconvex domains*, Indiana Univ. Math. J. **37** (1988), 145–163.
- [20] M. J. Martín, D. Vukotić, *Isometries of the Bloch space among the composition operators*, Bull. London Math. Soc., **39** (2007), 151–155.
- [21] W. Rudin, *Function Theory in Unit Ball of $\mathbb{C}^n$* , Springer–Verlag, New York, 1980.
- [22] W. Rudin, *Function Theory in Polydiscs*, W.A. Benjamin, Inc., New York, 1969.
- [23] R. M. Timoney, *Bloch functions in several complex variables, I*, Bull. London Math. Soc. **12** (1980), 241–267.
- [24] R. M. Timoney, *Bloch functions in several complex variables, II*, J. Reine Angew. Math. **319** (1980), 1–22.
- [25] J. H. Shi, L. Luo, *Composition operators on the Bloch space of several complex variables*, Acta Math. Sinica (N.S.) **16** (2000), 85–98.
- [26] G. Zhang, *Bloch constants of bounded symmetric domains*, Trans. Amer. Math. Soc. **349**(1997), 2941–2949.
- [27] K. Zhu, *Operator Theory in Function Spaces*, Marcel Dekker, New York, 1990.
- [28] K. Zhu, *Spaces of Holomorphic Functions in the Unit Ball*, Springer, New York, 2005.
- [29] Z. Zhou, J. Shi, *Composition operators on the Bloch space in polydisks*, Complex Variables **46** (2001), 73–88.
- [30] C. Xiong, *Norm of composition operators on the Bloch space*, Bull. Austral. Math. Soc. **70** (2004), 293–299.

GEORGE MASON UNIVERSITY, DEPARTMENT OF MATHEMATICS SCIENCES, FAIRFAX, VA 22030
E-mail address: `rallen2@gmu.edu`

GEORGE MASON UNIVERSITY, DEPARTMENT OF MATHEMATICS SCIENCES, FAIRFAX, VA 22030
E-mail address: `fcolonna@gmu.edu`